



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07D 211/42, 405/06, 401/12, 405/12, 405/14, A61K 31/445</b>		A1	(11) International Publication Number: <b>WO 97/30031</b> (43) International Publication Date: <b>21 August 1997 (21.08.97)</b>
(21) International Application Number: <b>PCT/EP97/00584</b>		(81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: <b>7 February 1997 (07.02.97)</b>			
(30) Priority Data: <b>96200379.4 15 February 1996 (15.02.96) EP</b> (34) Countries for which the regional or international application was filed: <b>DE et al.</b>			
(71) Applicant (for all designated States except US): <b>JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).</b>		Published <i>With international search report.</i>	
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>BOSMANS, Jean-Paul, R., M., A. [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). LOVE, Christopher, J. [GB/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). DECLEYN, Michel, A., J. [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). D'HAEN, Henri, E., F. [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).</b>			

## (54) Title: ESTERS OF 3-HYDROXY-PIPERIDINEMETHANOL DERIVATIVES

## (57) Abstract

The present invention of compounds of formula (I), a stereochemically isomeric form thereof, an N-oxide form thereof or a pharmaceutically acceptable acid addition salt thereof, R<sup>1</sup> is C<sub>1</sub>-alkyloxy, C<sub>2</sub>-alkenyloxy or C<sub>2</sub>-alkynyoxy; R<sup>2</sup> is hydrogen or C<sub>1</sub>-alkyloxy, or when taken together R<sup>1</sup> and R<sup>2</sup> may form a bivalent radical of formula wherein in said bivalent radicals one or two hydrogen atoms may be substituted with C<sub>1</sub>-alkyl; R<sup>3</sup> is hydrogen or halo; R<sup>4</sup> is hydrogen or C<sub>1</sub>-alkyl; L is C<sub>3</sub>-cycloalkyl, C<sub>5</sub>-cycloalkanone, C<sub>2</sub>-alkenyl optionally substituted with aryl, or L is a radical of formula -Alk-R<sup>5</sup>, Alk-X-R<sup>6</sup>, -Alk-Y-C(=O)-R<sup>8</sup>, or -Alk-Y-C(=O)-NR<sup>10</sup>R<sup>11</sup> wherein each Alk is C<sub>1</sub>-2alkanediyl; and R<sup>5</sup> is hydrogen, cyano, C<sub>1</sub>-alkylsulfonylaminoo, C<sub>3</sub>-cycloalkyl, C<sub>5</sub>-cycloalkanone, aryl, di(aryl)methyl or a heterocyclic ringsystem; R<sup>6</sup> is hydrogen, C<sub>1</sub>-alkyl, hydroxyC<sub>1</sub>-alkyl, C<sub>6</sub>-cycloalkyl, aryl or heterocyclic ringsystem; X is O, S, SO<sub>2</sub> or NR<sup>7</sup>; said R<sup>7</sup> being hydrogen, C<sub>1</sub>-alkyl or aryl; R<sup>8</sup> is hydrogen, C<sub>1</sub>-alkyl, C<sub>3</sub>-cycloalkyl, aryl, arylC<sub>1</sub>-alkyl, di(aryl)methyl, C<sub>1</sub>-alkyloxy or hydroxy; Y is NR<sup>9</sup> or a direct bond; said R<sup>9</sup> being hydrogen, C<sub>1</sub>-alkyl or aryl; R<sup>10</sup> and R<sup>11</sup> each independently are hydrogen, C<sub>1</sub>-alkyl, C<sub>3</sub>-cycloalkyl, aryl or arylC<sub>1</sub>-alkyl, or R<sup>10</sup> and R<sup>11</sup> combined with the nitrogen atom bearing R<sup>10</sup> and R<sup>11</sup> may form a pyrrolidinyl or piperidinyl ring both being optionally substituted with C<sub>1</sub>-alkyl, amino or mono or di(C<sub>1</sub>-alkyl)amino, or said R<sup>10</sup> and R<sup>11</sup> combined with the nitrogen bearing R<sup>10</sup> and R<sup>11</sup> may form a piperazine or a morpholine radical both being optionally substituted with C<sub>1</sub>-alkyl. Processes for preparing said products, formulations comprising said products and their use as a medicine are disclosed, in particular for treating conditions which are related to unpairment of gastric empaving.

